EVIDENCE SUMMARY: THE BENEFITS OF RELATIONAL CONTINUITY IN PRIMARY CARE

PREVENTIVE CARE 8/10

> studies showed improvements in preventive care

TILIZATION 61/64

studies showed positive results in lower utilization and hospitalization

MORTALITY 11/11

Ŵ

2017 UPDATE

studies showed reduced mortality

HEALTH 13/16

studies showed positive results in improved health

COST SAVINGS 16/17

\$

studies demonstrated _____cost savings

CARE QUALITY 10/16

studies showed positive results in overall care quality

SATISFACTION 15/16

studies showed increased patient satisfaction

ADHERENCE 6/6

studies showed improved self-management and treatment adherence

Evidence Summary - the benefits of relational continuity in primary care

October 2017

Question

What is the value of relational continuity with a primary health care provider?

Summary of findings

- A growing body of evidence points to the association of relational continuity with a single provider, and to a lesser extent, practice continuity with improved outcomes & satisfaction.
- For some measures, (utilization, mortality) the greater the degree of attachment, the better the outcomes.
- See below for overall outcomes and themes
- The numbers correspond with the numbered reference list below the summary
- This 2017 update includes 25 new articles, a new section on the value of relational continuity to patients and providers, and more detail on utilization outcomes and on the impact of continuity on complex patients.
- There is some variability in study quality, effect size, and in how relational continuity is defined. Study quality has not been evaluated in this summary.
- Please note **bolded** results reflect review articles that analyze multiple studies and may include peer reviewed articles and/or reports. As such, individual articles may be represented more than once.
- Should you require further clarification, please email: top@topalbertadoctors.org.

Relational continuity has been associated with:

- Fewer emergency visits & hospitalizations
 - 3, 4, 6, 7, 11, 17, 19, 22, 23, 28, 33, 35, 37, 38, 39, 42, 43, 46 (practice), 51, 53, 54, 55, 57, 59, 60, 61, 62, 65, 67, 72, 73, 77, 79, 80, 81, 82, 85, 86, 93, 95, 96, 102, 103, 105, 106, 107, 110, 113, 114, 116, 119, 120, 122(practice), 125, 127, 129, 131, 132, 136, 137
 - Several studies specifically found fewer ER visits & hospitalizations in patients with complex needs or a chronic disease.
 - 4, 22, 23, 28, 33, 37, 38, 53, 55, 59, 67, 73, 80, 81, 96, 105, 120, 122(practice), 129, 132

- Some studies found more of an impact on emergency visits for complex needs patients, compared to those without complex needs. 136
- o Greater degree of attachment was associated with a larger impact on this outcome
 - 3, 7, 11, 17, 22, 23, 28, 33, 51, 77, 122(practice), 125(practice vs provider), 127, 131, 137
- No significant impact on emergency visits & hospitalizations
 - 78 (Authors point to measurement issues. Still advocate for continuity), 112 (children)
- Increased emergency visits & hospitalizations
 - o 111 (patient surveys)
- Reduced specialty care utilization
 - o 32, 54
- Cost savings
 - 6, 12, 16(practice), 30 (practice), 37, 43, 46 (practice), 47, 55, 59, 65, 66, 83 (practice), 116, 122(practice), 129
 - Studies that found cost savings in patients with complex needs, or a chronic disease.
 30 (practice), 37, 55, 59, 122(practice)
 - Some studies found more of an impact on cost savings for complex needs or chronic disease patients, compared to those without complex needs. 122(practice)
- No significant cost savings
 - o **18**
- Reduced mortality
 - o 36, 53, 58, 62, 66, 71, 77, 87, 108, 109, 123
 - o The greater the degree of attachment, the greater the impact on mortality. 77
 - o Studies that found reduced mortality in patients with complex needs, or a chronic disease. 53, 71, 123
- Improved health & quality of life
 - o 3, 16(practice), 47, 52, 59, 66, 90 (practice), 101, 104, 108, 115, 117, 130
- No impact on health or quality of life
 - o 14, 20, 91
- Improved self-management and treatment adherence by patients
 - o 52, 92, 94, 99, 100, 133
- Improved preventive service delivery rates
 - o 10, 19, 29, **45**, 56, 75, 98, 128
- No significant impact on preventive services
 - o 18,44
- Improved quality of care
 - o 5, 34, **41**, 49, 59, 69, 76 (provider & practice), 116, 134, 135
 - o Studies that found improved quality in patients with complex needs, or a chronic disease. 59

- Some studies found more of an impact for patients with complex needs or a chronic disease, compared to those without complex needs. 135
- No significant/variable impact on quality of care
 - o 13(diabetes), 14(departing pcps), 18, 20 (cv risk factors), 84 (cancer), 94 (possible overprescribing)
- A decrease in overuse of medical procedures
 - o 24, 97
- Better patient-physician communication
 - o 68
- Improved patient satisfaction & experience
 - o 1, 8 (increased trust), 15, 31, **39, 40** (Some variable results though), 48, **57**(especially chronic disease patients), 63, 64, 70, 88, 89, 118, 121
- No impact on patient satisfaction & experience
 - o 18
- Do patients value continuity?
 - Yes, especially vulnerable patients, and those with complex conditions. 9, 15, 25, 48, 50, 70, 118, 121, 126
 - o Yes, but not patients with complex conditions 124
 - Some younger healthy patients or those with acute issues do not see the value 50, 126
- Improved staff satisfaction & experience
 - o 1, 2, 25, 26
- Do providers value continuity?
 - o Yes, especially for patients with complex conditions 2, 9, 21, 25, 26, 27, 121
- Higher performing primary practice
 - o 41, 74

Key articles to cite:

- Major studies & reviews
 - o 36, **39**, **50**, 58,59, 60, 64, **67**, 83, 86, 105, 121, 127
- Alberta results
 - o 60, 86, 121

Reference List

1985

1. Linn LS, Brook RH, Clark VA, Davies AR, Fink A, Kosecoff J. Physician and patient satisfaction as factors related to the organization of internal medicine group practices. Med Care. 1985 Oct;23(10):1171–8.

2. Blankfield RP, Kelly RB, Alemagno SA, King CM. Continuity of care in a family practice residency program. Impact on physician satisfaction. J Fam Pract. 1990 Jul;31(1):69–73.

1998

- 3. Mainous AG 3rd, Gill JM. The importance of continuity of care in the likelihood of future hospitalization: is site of care equivalent to a primary clinician? Am J Public Health. 1998 Oct;88(10):1539–41. <u>http://www.ncbi.nlm.nih.gov/pmc/articles/PMC1508474/</u>
- 4. Gill JM, Mainous AG 3rd. The role of provider continuity in preventing hospitalizations. Arch Fam Med. 1998 Aug;7(4):352–7. <u>http://triggered.stanford.clockss.org/ServeContent?url=http%3A%2F%2Farchfami.ama-assn.org%2Fcgi%2Fcontent%2Ffull%2F7%2F4%2F352</u>
- O'Connor PJ, Desai J, Rush WA, Cherney LM, Solberg LI, Bishop DB. Is having a regular provider of diabetes care related to intensity of care and glycemic control? J Fam Pract. 1998 Oct;47(4):290–7.

1999

6. Raddish M, Horn SD, Sharkey PD. Continuity of care: is it cost effective? Am J Manag Care. 1999 Jun;5(6):727–34.

2001

- 7. Christakis DA, Mell L, Koepsell TD, Zimmerman FJ, Connell FA. Association of lower continuity of care with greater risk of emergency department use and hospitalization in children. Pediatrics. 2001 Mar;107(3):524–9.
- Mainous AG, Baker R, Love MM, Gray DP, Gill JM. Continuity of care and trust in one's physician: evidence from primary care in the United States and the United Kingdom. Fam Med. 2001 Jan;33(1):22–7.
- 9. Kearley KE, Freeman GK, Heath A. An exploration of the value of the personal doctor-patient relationship in general practice. Br J Gen Pract. 2001 Sep;51(470):712–8. https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1314098/

2002

 Xu KT. Usual source of care in preventive service use: a regular doctor versus a regular site. Health Serv Res. 2002 Dec;37(6):1509–29. <u>http://www.ncbi.nlm.nih.gov/pmc/articles/PMC1464041/</u>

2003

11. Burge F, Lawson B, Johnston G. Family physician continuity of care and emergency department use in end-of-life cancer care. Med Care. 2003 Aug;41(8):992–1001.

- 12. De Maeseneer JM, De Prins L, Gosset C, Heyerick J. Provider continuity in family medicine: does it make a difference for total health care costs? Ann Fam Med. 2003 Oct;1(3):144–8. <u>http://www.ncbi.nlm.nih.gov/pmc/articles/PMC1466579/</u>
- 13. Gill JM, Mainous AG, Diamond JJ, Lenhard MJ. Impact of provider continuity on quality of care for persons with diabetes mellitus. Ann Fam Med. 2003 Sep;1(3):162–70. http://www.ncbi.nlm.nih.gov/pmc/articles/PMC1466582
- 14. Pereira AG, Kleinman KP, Pearson SD. Leaving the practice: effects of primary care physician departure on patient care. Arch Intern Med. 2003 Dec 8;163(22):2733–6. http://archinte.jamanetwork.com/article.aspx?doi=10.1001/archinte.163.22.2733
- 15. Nutting PA, Goodwin MA, Flocke SA, Zyzanski SJ, Stange KC. Continuity of primary care: to whom does it matter and when? Ann Fam Med. 2003 Oct;1(3):149–55. https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1466596/

- 16. Starfield B, Shi L. The medical home, access to care, and insurance: a review of evidence. Pediatrics. 2004 May 1;113(Supplement 4):1493–8.
- Brousseau DC, Meurer JR, Isenberg ML, Kuhn EM, Gorelick MH. Association between infant continuity of care and pediatric emergency department utilization. Pediatrics. 2004 Apr;113(4):738–41. <u>http://pediatrics.aappublications.org/content/113/4/738.long</u>
- Parkerton PH, Smith DG, Straley HL. Primary care practice coordination versus physician continuity. Fam Med. 2004 Jan;36(1):15–21. http://www.stfm.org/fmhub/fm2004/January/Patricia15.pdf

2005

- Saultz JW, Lochner J. Interpersonal continuity of care and care outcomes: a critical review. Ann Fam Med. 2005 Mar;3(2):159–66. http://www.ncbi.nlm.nih.gov/pmc/articles/PMC1466859/
- 20. Litaker D, Ritter C, Ober S, Aron D. Continuity of care and cardiovascular risk factor management: does care by a single clinician add to informational continuity provided by electronic medical records? Am J Manag Care. 2005 Nov;11(11):689–96. <u>http://www.ajmc.com/publications/issue/2005/2005-11-vol11-n11/Nov05-2207p689-696/</u>
- Stokes T, Tarrant C, Mainous AG, Schers H, Freeman G, Baker R. Continuity of Care: Is the Personal Doctor Still Important? A Survey of General Practitioners and Family Physicians in England and Wales, the United States, and the Netherlands. Ann Fam Med. 2005 Jul 1;3(4):353–9. http://www.annfammed.org/content/3/4/353.full

- 22. Cree M, Bell NR, Johnson D, Carriere KC. Increased continuity of care associated with decreased hospital care and emergency department visits for patients with asthma. Dis Manag. 2006 Feb;9(1):63–71.
- 23. Menec VH, Sirski M, Attawar D, Katz A. Does continuity of care with a family physician reduce hospitalizations among older adults? J Health Serv Res Policy. 2006 Oct 1;11(4):196–201. http://hsr.sagepub.com/content/11/4/196.long

- 24. van Walraven C, Cernat G, Austin PC. Effect of provider continuity on test repetition. Clin Chem. 2006 Dec;52(12):2219–28. <u>http://www.clinchem.org/content/52/12/2219.long</u>
- 25. Guthrie B, Wyke S. Personal continuity and access in UK general practice: a qualitative study of general practitioners' and patients' perceptions of when and how they matter. BMC Family Practice. 2006 Feb 24;7:11. <u>https://bmcfampract.biomedcentral.com/articles/10.1186/1471-2296-7-11</u>
- 26. Ridd M, Shaw A, Salisbury C. 'Two sides of the coin'--the value of personal continuity to GPs: a qualitative interview study. Fam Pract. 2006 Aug;23(4):461–8.
- 27. Schers H, Bor H, van den Bosch W, Grol R. GPs' attitudes to personal continuity: findings from everyday practice differ from postal surveys. Br J Gen Pract. 2006 Jul 1;56(528):536–8. https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1872065/

 Ionescu-Ittu R, McCusker J, Ciampi A, Vadeboncoeur A-M, Roberge D, Larouche D, et al. Continuity of primary care and emergency department utilization among elderly people. CMAJ. 2007 Nov 20;177(11):1362–8. <u>http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2072991/</u>

2008

29. Blewett LA, Johnson PJ, Lee B, Scal PB. When a usual source of care and usual provider matter: adult prevention and screening services. J Gen Intern Med. 2008 Sep;23(9):1354–60. http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2518015/

2009

- Hollander MJ, Kadlec H, Hamdi R, Tessaro A. Increasing value for money in the Canadian healthcare system: new findings on the contribution of primary care services. Healthc Q. 2009;12(4):32–44.
- 31. Scobie A, MacKinnon NJ, Higgins S, Etchegary H, Church R. The medical home in Canada: patient perceptions of quality and safety. Healthc Manage Forum. 2009;22(1):47–51.
- 32. Starfield B, Chang H-Y, Lemke KW, Weiner JP. Ambulatory specialist use by nonhospitalized patients in us health plans: correlates and consequences. J Ambul Care Manage. 2009 Sep;32(3):216–25.
- 33. Knight JC, Dowden JJ, Worrall GJ, Gadag VG, Murphy MM. Does higher continuity of family physician care reduce hospitalizations in elderly people with diabetes? Popul Health Manag. 2009 Apr;12(2):81–6.
- 34. Atlas SJ, Grant RW, Ferris TG, Chang Y, Barry MJ. Patient-physician connectedness and quality of primary care. Ann Intern Med. 2009 Mar 3;150(5):325–35. http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2975389/

2010

35. Cheng S-H, Chen C-C, Hou Y-F. A longitudinal examination of continuity of care and avoidable hospitalization: evidence from a universal coverage health care system. Arch Intern Med.

2010 Oct 11;170(18):1671-7.

http://archinte.jamanetwork.com/article.aspx?doi=10.1001/archinternmed.2010.340

- 36. Wolinsky FD, Bentler SE, Liu L, Geweke JF, Cook EA, Obrizan M, et al. Continuity of care with a primary care physician and mortality in older adults. J Gerontol A Biol Sci Med Sci. 2010 Apr;65(4):421–8. <u>http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2844057/</u>
- Hong JS, Kang HC, Kim J. Continuity of care for elderly patients with diabetes mellitus, hypertension, asthma, and chronic obstructive pulmonary disease in Korea. J Korean Med Sci. 2010 Sep;25(9):1259–71. <u>http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2923791/</u>
- 38. Lin W, Huang I-C, Wang S-L, Yang M-C, Yaung C-L. Continuity of diabetes care is associated with avoidable hospitalizations: evidence from Taiwan's National Health Insurance scheme. Int J Qual Health Care. 2010 Feb;22(1):3–8. <u>http://intqhc.oxfordjournals.org/content/22/1/3.long</u>
- 39. van Walraven C, Oake N, Jennings A, Forster AJ. The association between continuity of care and outcomes: a systematic and critical review. J Eval Clin Pract. 2010 Oct;16(5):947–56.
- Adler R, Vasiliadis A, Bickell N. The relationship between continuity and patient satisfaction: a systematic review. Fam Pract. 2010 Apr;27(2):171–8. http://fampra.oxfordjournals.org/content/27/2/171.long
- 41. Kringos DS, Boerma WG, Hutchinson A, van der Zee J, Groenewegen PP. The breadth of primary care: a systematic literature review of its core dimensions. BMC Health Services Research. 2010;10:65. <u>http://bmchealthservres.biomedcentral.com/articles/10.1186/1472-6963-10-65</u>

2011

- 42. Cheng S-H, Hou Y-F, Chen C-C. Does continuity of care matter in a health care system that lacks referral arrangements? Health Policy Plan. 2011 Mar;26(2):157–62. http://heapol.oxfordjournals.org/content/26/2/157.long
- Chen C-C, Chen S-H. Better continuity of care reduces costs for diabetic patients. Am J Manag Care. 2011 Jun;17(6):420–7. <u>http://www.ajmc.com/publications/issue/2011/2011-6-vol17-n6/AJMC_11jun_ChenChen_420to27/</u>
- 44. McRae I, Yen L, Gillespie J, Douglas K. Patient affiliation with GPs in Australia--who is and who is not and does it matter? Health Policy. 2011 Nov;103(1):16–23.

- Kim MY, Kim JH, Choi I-K, Hwang IH, Kim SY. Effects of having usual source of care on preventive services and chronic disease control: a systematic review. Korean J Fam Med. 2012 Nov;33(6):336–45. <u>http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3526716/</u>
- 46. Anderson LH, Flottemesch TJ, Fontaine P, Solberg LI, Asche SE. Patient medical group continuity and healthcare utilization. Am J Manag Care. 2012 Aug;18(8):450–7. <u>http://www.ajmc.com/publications/issue/2012/2012-8-vol18-n8/Patient-Medical-Group-Continuity-and-Healthcare-Utilization/</u>
- 47. Chu H-Y, Chen C-C, Cheng S-H. Continuity of care, potentially inappropriate medication, and health care outcomes among the elderly: evidence from a longitudinal analysis in Taiwan. Med Care. 2012 Nov;50(11):1002–9.

- 48. Crooks VA, Agarwal G, Harrison A. Chronically ill Canadians' experiences of being unattached to a family doctor: a qualitative study of marginalized patients in British Columbia. BMC Family Practice. 2012 Jul 16;13(1):69. <u>http://www.biomedcentral.com/1471-2296/13/69</u>
- Devoe JE, Tillotson CJ, Wallace LS, Lesko SE, Angier H. The effects of health insurance and a usual source of care on a child's receipt of health care. J Pediatr Health Care. 2012 Oct;26(5):e25–35. <u>http://www.ncbi.nlm.nih.gov/pmc/articles/pmid/22920780/</u>
- 50. Waibel S, Henao D, Aller M-B, Vargas I, Vázquez M-L. What do we know about patients' perceptions of continuity of care? A meta-synthesis of qualitative studies. Int J Qual Health Care. 2012 Feb;24(1):39–48.

http://intqhc.oxfordjournals.org/cgi/pmidlookup?view=long&pmid=22146566

2013

- 51. Nyweide DJ, Anthony DL, Bynum JW, et al. Continuity of care and the risk of preventable hospitalization in older adults. JAMA Intern Med. 2013 Sep 16; <u>http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3877937/</u>
- 52. Chen C-C, Tseng C-H, Cheng S-H. Continuity of care, medication adherence, and health care outcomes among patients with newly diagnosed type 2 diabetes: a longitudinal analysis. Med Care. 2013 Mar;51(3):231–7.
- McAlister FA, Youngson E, Bakal JA, Kaul P, Ezekowitz J, van Walraven C. Impact of physician continuity on death or urgent readmission after discharge among patients with heart failure. CMAJ. 2013 Oct 1;185(14):E681–9. http://www.ncbi.nlm.nih.gov/pmc/articles/pmid/23959284/
- 54. Hansen AH, Halvorsen PA, Aaraas IJ, Førde OH. Continuity of GP care is related to reduced specialist healthcare use: a cross-sectional survey. Br J Gen Pract. 2013 Jul;63(612):482–9. http://www.ncbi.nlm.nih.gov/pmc/articles/pmid/23834885/
- 55. Hong J-S, Kang H-C. Continuity of ambulatory care and health outcomes in adult patients with type 2 diabetes in Korea. Health Policy. 2013 Feb;109(2):158–65.
- 56. Schoueri-Mychasiw N, McDonald PW. Factors associated with underscreening for cervical cancer among women in Canada. Asian Pac J Cancer Prev. 2013;14(11):6445–50. http://journal.waocp.org/article 28316 30e54434bd694c08a67f2ad226281157.pdf
- 57. Health Quality Ontario. Continuity of care to optimize chronic disease management in the community setting: an evidence-based analysis. Ont Health Technol Assess Ser. 2013;13(6):1–41.http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3806147/
- 58. Leleu H, Minvielle E. Relationship between longitudinal continuity of primary care and likelihood of death: analysis of national insurance data. PLoS ONE. 2013;8(8):e71669. <u>http://www.ncbi.nlm.nih.gov/pmc/articles/pmid/23990970/</u>

- 59. Hussey PS, Schneider EC, Rudin RS, Fox DS, Lai J, Pollack CE. Continuity and the costs of care for chronic disease. JAMA Intern Med. 2014 May;174(5):742–8. https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4075052/
- 60. Health Quality Council of Alberta. Primary care measurement initiative report [Internet]. Calgary, AB: HQCA; 2014 Mar. Available from: <u>http://hqca.ca/news/2014/03/primary-care-measurement-initiative-report-released/</u>

- 61. Chaiyachati KH, Gordon K, Long T, Levin W, Khan A, Meyer E, et al. Continuity in a VA patientcentered medical home reduces emergency department visits. PLoS ONE. 2014;9(5):e96356. <u>http://www.ncbi.nlm.nih.gov/pubmed/24867300</u>
- 62. Nelson K, Sun H, Dolan E, Maynard C, Beste L, Bryson C, et al. Elements of the patientcentered medical home associated with health outcomes among veterans: the role of primary care continuity, expanded access, and care coordination. J Ambul Care Manage. 2014 Dec;37(4):331–8.
- 63. Wetmore S, Boisvert L, Graham E, Hall S, Hartley T, Wright L, et al. Patient satisfaction with access and continuity of care in a multidisciplinary academic family medicine clinic. Can Fam Physician. 2014 Apr;60(4):e230–6. http://www.ncbi.nlm.nih.gov/pmc/articles/pmid/24733343/
- 64. Rhodes P, Sanders C, Campbell S. Relationship continuity: when and why do primary care patients think it is safer? Br J Gen Pract. 2014 Dec;64(629):e758–64. http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4240148/
- 65. Romaire MA, Haber SG, Wensky SG, McCall N. Primary care and specialty providers: an assessment of continuity of care, utilization, and expenditures. Med Care. 2014 Dec;52(12):1042–9.
- 66. Shin DW, Cho J, Yang HK, Park JH, Lee H, Kim H, et al. Impact of continuity of care on mortality and health care costs: a nationwide cohort study in Korea. Ann Fam Med. 2014 Dec;12(6):534–41. <u>http://www.ncbi.nlm.nih.gov/pmc/articles/pmid/25384815/</u>
- 67. Van Loenen T, van den Berg MJ, Westert GP, Faber MJ. Organizational aspects of primary care related to avoidable hospitalization: a systematic review. Fam Pract. 2014 Oct;31(5):502–16. <u>https://academic.oup.com/fampra/article-lookup/doi/10.1093/fampra/cmu053</u>
- 68. Katz DA, McCoy K, Sarrazin MV. Does improved continuity of primary care affect clinicianpatient communication in VA? J Gen Intern Med. 2014 Jul;29 Suppl 2:S682–8. <u>http://www.ncbi.nlm.nih.gov/pmc/articles/pmid/24072718/</u>
- 69. Cook JA, Burke-Miller JK, Grey DD, Cocohoba J, Liu C, Schwartz RM, et al. Do HIV-positive women receive depression treatment that meets best practice guidelines? AIDS Behav. 2014 Jun;18(6):1094–102. <u>http://www.ncbi.nlm.nih.gov/pmc/articles/pmid/24402689/</u>
- Al-Azri M, Al-Ramadhani R, Al-Rawahi N, Al-Shafee K, Al-Hinai M, Al-Maniri A. Patients' attitudes and experiences of relational continuity in semi-urban general practices in Oman. Fam Pract. 2014 Jun;31(3):303–10. <u>https://academic.oup.com/fampra/article-lookup/doi/10.1093/fampra/cmu004</u>
- Spatz ES, Sheth SD, Gosch KL, Desai MM, Spertus JA, Krumholz HM, et al. Usual source of care and outcomes following acute myocardial infarction. J Gen Intern Med. 2014 Jun;29(6):862– 9. <u>http://www.ncbi.nlm.nih.gov/pmc/articles/pmid/24553957/</u>
- Huntley A, Lasserson D, Wye L, Morris R, Checkland K, England H, et al. Which features of primary care affect unscheduled secondary care use? A systematic review. BMJ Open. 2014;4(5):e004746. <u>http://bmjopen.bmj.com/content/4/5/e004746.full</u>
- Maciejewski ML, Powers BJ, Sanders LL, Farley JF, Hansen RA, Sleath B, et al. The intersection of patient complexity, prescriber continuity and acute care utilization. J Gen Intern Med. 2014 Apr;29(4):594–601. <u>http://www.ncbi.nlm.nih.gov/pmc/articles/pmid/24408277/</u>
- 74. Bodenheimer T, Ghorob A, Willard-Grace R, Grumbach K. The 10 building blocks of highperforming primary care. Ann Fam Med. 2014 Apr;12(2):166–71. <u>http://www.ncbi.nlm.nih.gov/pmc/articles/pmid/24615313/</u>
- 75. Brown ML, Klabunde CN, Cronin KA, White MC, Richardson LC, McNeel TS. Challenges in meeting Healthy People 2020 objectives for cancer-related preventive services, National

Health Interview Survey, 2008 and 2010. Prev Chronic Dis. 2014;11:E29. https://www.cdc.gov/pcd/issues/2014/13_0174.htm

- 76. Cheng S-H, Chen C-C. Effects of continuity of care on medication duplication among the elderly. Med Care. 2014 Feb;52(2):149–56.
- 77. Almaawiy U, Pond GR, Sussman J, Brazil K, Seow H. Are family physician visits and continuity of care associated with acute care use at end-of-life? A population-based cohort study of homecare cancer patients. Palliat Med. 2014 Feb;28(2):176–83.
- 78. Bentler SE, Morgan RO, Virnig BA, Wolinsky FD. The association of longitudinal and interpersonal continuity of care with emergency department use, hospitalization, and mortality among Medicare beneficiaries. PLoS ONE. 2014;9(12):e115088. <u>http://www.ncbi.nlm.nih.gov/pmc/articles/pmid/25531108/</u>

2015

- 79. Cho KH, Lee SG, Jun B, Jung B-Y, Kim J-H, Park E-C. Effects of continuity of care on hospital admission in patients with type 2 diabetes: analysis of nationwide insurance data. BMC Health Serv Res. 2015;15:107. <u>http://www.ncbi.nlm.nih.gov/pmc/articles/pmid/25879858/</u>
- Bayliss EA, Ellis JL, Shoup JA, Zeng C, McQuillan DB, Steiner JF. Effect of continuity of care on hospital utilization for seniors with multiple medical conditions in an integrated health care system. Ann Fam Med. 2015 Mar;13(2):123–9. <u>http://www.ncbi.nlm.nih.gov/pmc/articles/pmid/25755033/</u>
- 81. Lin I-P, Wu S-C, Huang S-T. Continuity of care and avoidable hospitalizations for chronic obstructive pulmonary disease (COPD). J Am Board Fam Med. 2015 Apr;28(2):222–30. http://www.jabfm.org/cgi/pmidlookup?view=long&pmid=25748763
- 82. Liao P-J, Lin Z-Y, Huang J-C, Hsu K-H. The relationship between type 2 diabetic patients' early medical care-seeking consistency to the same clinician and health care system and their clinical outcomes. Medicine (Baltimore). 2015 Feb;94(7):e554. http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4554162/
- 83. Hollander MJ, Kadlec H. Financial implications of the continuity of primary care. Perm J. 2015;19(1):4–10. <u>http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4315369/</u>
- Ridd MJ, Ferreira DLS, Montgomery AA, Salisbury C, Hamilton W. Patient-doctor continuity and diagnosis of cancer: electronic medical records study in general practice. Br J Gen Pract. 2015 May;65(634):e305–11. <u>http://www.ncbi.nlm.nih.gov/pmc/articles/pmid/25918335/</u>
- 85. Comino EJ, Islam MF, Tran DT, Jorm L, Flack J, Jalaludin B, et al. Association of processes of primary care and hospitalisation for people with diabetes: A record linkage study. Diabetes Res Clin Pract. 2015 May;108(2):296–305.
- 86. Canadian Institute for Health Information. Continuity of care with family medicine physicians: why it matters. CIHI; 2015. <u>https://secure.cihi.ca/estore/productSeries.htm?pc=PCC1299</u>
- Cho KH, Kim YS, Nam CM, Kim TH, Kim SJ, Han K-T, et al. The Association between Continuity of Care and All-Cause Mortality in Patients with Newly Diagnosed Obstructive Pulmonary Disease: A Population-Based Retrospective Cohort Study, 2005-2012. PLoS ONE. 2015;10(11):e0141465.

http://journals.plos.org/plosone/article?id=10.1371/journal.pone.0141465

88. Du Z, Liao Y, Chen C-C, Hao Y, Hu R. Usual source of care and the quality of primary care: a survey of patients in Guangdong province, China. Int J Equity Health. 2015;14:60. <u>https://equityhealthj.biomedcentral.com/articles/10.1186/s12939-015-0189-4</u>

- 89. Finney Rutten LJ, Agunwamba AA, Beckjord E, Hesse BW, Moser RP, Arora NK. The Relation Between Having a Usual Source of Care and Ratings of Care Quality: Does Patient-Centered Communication Play a Role? J Health Commun. 2015;20(7):759–65.
- 90. Gong Y-H, Yoon S-J, Seo H, Kim D. Associations Between the Continuity of Ambulatory Care of Adult Diabetes Patients in Korea and the Incidence of Macrovascular Complications. J Prev Med Public Health. 2015 Jul;48(4):188–94. http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4542297/
- 91. Hanafi NS, Abdullah A, Lee PY, Liew SM, Chia YC, Khoo EM. Personal Continuity of Care in a University-Based Primary Care Practice: Impact on Blood Pressure Control. PLoS ONE. 2015;10(7):e0134030. <u>http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4516235/</u>
- 92. Hansen RA, Voils CI, Farley JF, Powers BJ, Sanders LL, Sleath B, et al. Prescriber continuity and medication adherence for complex patients. Ann Pharmacother. 2015 Mar;49(3):293–302.
- 93. Katz DA, McCoy KD, Vaughan-Sarrazin MS. Does Greater Continuity of Veterans Administration Primary Care Reduce Emergency Department Visits and Hospitalization in Older Veterans? J Am Geriatr Soc. 2015 Dec;63(12):2510–8. <u>http://onlinelibrary.wiley.com/doi/10.1111/jgs.13841/full</u>
- 94. Lam TP, Wun YT, Lam KF, Sun KS. Differences in antibiotic use between patients with and without a regular doctor in Hong Kong. BMC Pharmacol Toxicol. 2015;16:40. <u>http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4681134/</u>
- 95. Pourat N, Davis AC, Chen X, Vrungos S, Kominski GF. In California, Primary Care Continuity Was Associated With Reduced Emergency Department Use And Fewer Hospitalizations. Health Aff (Millwood). 2015 Jul;34(7):1113–20.
- 96. Raphael JL, Cooley WC, Vega A, Kowalkowski MA, Tran X, Treadwell J, et al. Outcomes for Children with Chronic Conditions Associated with Parent- and Provider-reported Measures of the Medical Home. J Health Care Poor Underserved. 2015 May;26(2):358–76. <u>http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4460004/</u>
- 97. Romano MJ, Segal JB, Pollack CE. The Association Between Continuity of Care and the Overuse of Medical Procedures. JAMA Intern Med. 2015 Jul;175(7):1148–54. http://archinte.jamanetwork.com/article.aspx?articleid=2294233
- 98. VanGompel ECW, Jerant AF, Franks PM. Primary Care Attributes Associated with Receipt of Preventive Care Services: A National Study. J Am Board Fam Med. 2015 Dec;28(6):733–41. <u>http://www.jabfm.org/content/28/6/733.full</u>
- 99. Warren JR, Falster MO, Tran B, Jorm L. Association of Continuity of Primary Care and Statin Adherence. PLoS ONE. 2015;10(10):e0140008. http://journals.plos.org/plosone/article?id=10.1371/journal.pone.0140008

- 100. Chen C-C, Cheng S-H. Continuity of care and changes in medication adherence among patients with newly diagnosed diabetes. Am J Manag Care. 2016 Feb;22(2):136–42. <u>http://www.ajmc.com/journals/issue/2016/2016-vol22-n2/Continuity-of-Care-and-Changes-in-Medication-Adherence-Among-Patients-With-Newly-Diagnosed-Diabetes/</u>
- 101. Chen H-M, Tu Y-H, Chen C-M. Effect of Continuity of Care on Quality of Life in Older Adults With Chronic Diseases: A Meta-Analysis. Clin Nurs Res. 2016 Jan 20;
- 102. Cho KH, Nam CM, Choi Y, Choi J-W, Lee S-H, Park E-C. Impact of continuity of care on preventable hospitalization of patients with type 2 diabetes: a nationwide Korean cohort study, 2002-10. Int J Qual Health Care. 2016 Jun 6;

- 103. Colligan EM, Pines JM, Colantuoni E, Wolff JL. Factors Associated With Frequent Emergency Department Use in the Medicare Population. Med Care Res Rev. 2016 Mar 29;
- 104. Dinkler JM, Sugar CA, Escarce JJ, Ong MK, Mangione CM. Does Age Matter? Association Between Usual Source of Care and Hypertension Control in the US Population: Data From NHANES 2007-2012. Am J Hypertens. 2016 Feb 16; https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5006109/
- 105. Gruneir A, Bronskill SE, Maxwell CJ, Bai YQ, Kone AJ, Thavorn K, et al. The association between multimorbidity and hospitalization is modified by individual demographics and physician continuity of care: a retrospective cohort study. BMC Health Serv Res. 2016;16(1):154. <u>https://bmchealthservres.biomedcentral.com/articles/10.1186/s12913-016-1415-5</u>
- 106. Hsu C-H, Chou Y-J, Pu C. The Effect of Continuity of Care on Emergency Room Use for Diabetic Patients Varies by Disease Severity. J Epidemiol. 2016 Feb 20; https://www.jstage.jst.go.jp/article/jea/advpub/0/advpub_JE20150045/_pdf
- 107. Huang S-T, Wu S-C, Hung Y-N, Lin I-P. Effects of continuity of care on emergency department utilization in children with asthma. Am J Manag Care. 2016 Jan;22(1):e31-37. http://www.ajmc.com/pubMed.php?pii=86485
- 108. Lustman A, Comaneshter D, Vinker S. Interpersonal continuity of care and type two diabetes. Prim Care Diabetes. 2016 Jun;10(3):165–70.
- 109. Maarsingh OR, Henry Y, van de Ven PM, Deeg DJ. Continuity of care in primary care and association with survival in older people: a 17-year prospective cohort study. Br J Gen Pract. 2016 Jun 20;
- 110. Nam YS, Cho KH, Kang H-C, Lee K-S, Park E-C. Greater continuity of care reduces hospital admissions in patients with hypertension: An analysis of nationwide health insurance data in Korea, 2011-2013. Health Policy. 2016 Jun;120(6):604–11.
- 111. Thanh NX, Rapoport J. Health services utilization of people having and not having a regular doctor in Canada. Int J Health Plann Manage. 2016 Feb 10;
- 112. Turbitt E, Freed GL. Regular source of primary care and emergency department use of children in Victoria. J Paediatr Child Health. 2016 Mar;52(3):303–7.
- 113. Van Loenen T, Faber MJ, Westert GP, Van den Berg MJ. The impact of primary care organization on avoidable hospital admissions for diabetes in 23 countries. Scand J Prim Health Care. 2016 Mar;34(1):5–12. <u>http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4911022/</u>
- 114. Vogt V, Koller D, Sundmacher L. Continuity of care in the ambulatory sector and hospital admissions among patients with heart failure in Germany. Eur J Public Health. 2016 Mar 9;
- 115. Weir DL, McAlister FA, Majumdar SR, Eurich DT. The Interplay Between Continuity of Care, Multimorbidity, and Adverse Events in Patients With Diabetes. Med Care. 2016 Apr;54(4):386–93.
- 116. Witt WP, Fullerton C, Chow C, Gokhale M, Naeger S, Walsh C, et al. Impact of Having a Usual Source of Care on Health Care Outcomes among Children with Serious Emotional Disturbance. Acad Pediatr. 2016 Jun 8;
- 117. Ye T, Sun X, Tang W, Miao Y, Zhang Y, Zhang L. Effect of continuity of care on healthrelated quality of life in adult patients with hypertension: a cohort study in China. BMC Health Serv Res. 2016 Nov 28;16(1):674.

https://www.ncbi.nlm.nih.gov/pmc/articles/pmid/27894298/

118. Reid J, Cormack D, Crowe M. The significance of relational continuity of care for Māori patient engagement with predominantly non-Māori doctors: findings from a qualitative study. Aust N Z J Public Health. 2016 Apr;40(2):120–5. http://onlinelibrary.wiley.com/doi/10.1111/1753-6405.12447/full

- 119. Pu C, Chou Y-J. The impact of continuity of care on emergency room use in a health care system without referral management: an instrumental variable approach. Ann Epidemiol. 2016 Mar;26(3):183–8. <u>http://www.annalsofepidemiology.org/article/S1047-2797(16)30004-7/fulltext</u>
- 120. Lai Y-R, Yang Y-S, Tsai M-L, Lu Y-L, Kornelius E, Huang C-N, et al. Impact of potentially inappropriate medication and continuity of care in a sample of Taiwan elderly patients with diabetes mellitus who have also experienced heart failure. Geriatr Gerontol Int. 2016 Oct;16(10):1117–26.
- 121. Health Quality Council of Alberta. Understanding patient and provider experiences with relationship, information, and management continuity. Calgary, AB: Health Quality Council of Alberta; 2016 Aug. Available from: <u>http://hqca.ca/studies-and-reviews/relationship-information-and-management-continuity/relationship-information-and-management-continuity/</u>

- 122. Sveréus S, Larsson K, Rehnberg C. Clinic continuity of care, clinical outcomes and direct costs for COPD in Sweden: a population based cohort study. Eur Clin Respir J. 2017;4(1):1290193. https://www.ncbi.nlm.nih.gov/pmc/articles/pmid/28326179/
- 123. Pan C-C, Kung P-T, Chiu L-T, Liao YP, Tsai W-C. Patients with diabetes in pay-forperformance programs have better physician continuity of care and survival. Am J Manag Care. 2017 Feb 1;23(2):e57–66. <u>http://www.ajmc.com/journals/issue/2017/2017-vol23n2/Patients-With-Diabetes-in-Pay-for-Performance-Programs-Have-Better-Physician-Continuity-of-Care-and-Survival/</u>
- 124. Mendes FRP, Gemito MLGP, Caldeira E do C, Serra I da C, Casas-Novas MV. Continuity of care from the perspective of users. Cien Saude Colet. 2017 Mar;22(3):841–53. <u>http://www.scielo.br/scielo.php?pid=S1413-</u> <u>81232017002300841&script=sci_arttext&tlng=en#B32</u>
- 125. Kim KM, Jeon H, Lee JH. Having a Physician Rather than a Place as a Usual Source of Care Would Be Better - from 2012 Korea Health Panel Data. J Korean Med Sci. 2017 Jan;32(1):4– 12. <u>https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5143297/</u>
- 126. Ehman KM, Deyo-Svendsen M, Merten Z, Kramlinger AM, Garrison GM. How Preferences for Continuity and Access Differ Between Multimorbidity and Healthy Patients in a Team Care Setting. J Prim Care Community Health. 2017 Apr 1;2150131917704556.
- 127. Barker I, Steventon A, Deeny SR. Association between continuity of care in general practice and hospital admissions for ambulatory care sensitive conditions: cross sectional study of routinely collected, person level data. BMJ. 2017 Feb 1;356:j84. http://www.bmj.com/content/356/bmj.j84
- 128. Arnold LD, McGilvray MM, Kyle Cooper J, James AS. Inadequate Cancer Screening: Lack of Provider Continuity is a Greater Obstacle than Medical Mistrust. J Health Care Poor Underserved. 2017;28(1):362–77. https://www.ncbi.nlm.nih.gov/pmc/articles/pmid/28239007/
- 129. Amjad H, Carmichael D, Austin AM, Chang C-H, Bynum JPW. Continuity of Care and Health Care Utilization in Older Adults With Dementia in Fee-for-Service Medicare. JAMA Intern Med. 2016 Sep 1;176(9):1371–8. https://www.ncbi.nlm.nih.gov/pmc/articles/pmid/27454945/

- 130. Maciejewski ML, Hammill BG, Bayliss EA, Ding L, Voils CI, Curtis LH, et al. Prescriber Continuity and Disease Control of Older Adults. Med Care. 2017 Apr;55(4):405–10.
- 131. Nyweide DJ, Bynum JPW. Relationship Between Continuity of Ambulatory Care and Risk of Emergency Department Episodes Among Older Adults. Ann Emerg Med. 2017 Apr;69(4):407–415.e3. <u>http://www.annemergmed.com/article/S0196-0644(16)30305-5/fulltext</u>
- 132. Kao Y-H, Wu S-C. Effect of Continuity of Care on Emergency Department Visits in Elderly Patients with Asthma in Taiwan. J Am Board Fam Med. 2017 Jun;30(3):384–95. <u>http://www.jabfm.org/cgi/pmidlookup?view=long&pmid=28484073</u>
- 133. Dossa AR, Moisan J, Guénette L, Lauzier S, Grégoire J-P. Association between interpersonal continuity of care and medication adherence in type 2 diabetes: an observational cohort study. CMAJ Open. 2017 May 8;5(2):E359–64. <u>http://cmajopen.ca/content/5/2/E359.full</u>
- 134. Keim-Malpass J, Lindley LC. End-of-Life Transitions and Hospice Utilization for Adolescents: Does Having a Usual Source of Care Matter? J Hosp Palliat Nurs. 2017 Aug;19(4):376–82.
- 135. Guo J-Y, Chou Y-J, Pu C. Effect of Continuity of Care on Drug-Drug Interactions. Med Care. 2017 Aug;55(8):744–51.
- 136. Enlow E, Passarella M, Lorch SA. Continuity of Care in Infancy and Early Childhood Health Outcomes. Pediatrics. 2017 Jul;140(1).
- 137. Kohnke H, Zielinski A. Association between continuity of care in Swedish primary care and emergency services utilisation: a population-based cross-sectional study. Scand J Prim Health Care. 2017 Jun;35(2):113–9. <u>https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5499310/</u>

More Information

Please contact the ACTT librarian (<u>actt@albertadoctors.org</u>) if you require full text access to any of the articles without links or for more information.